

## LETTER

# Ecological genomics predicts climate vulnerability in an endangered southwestern songbird

Kristen Ruegg,<sup>1,2\*†</sup> Rachael A. Bay,<sup>1,3,4†</sup> Eric C. Anderson,<sup>3,4</sup> James F. Saracco,<sup>5</sup> Ryan J. Harrigan,<sup>1</sup> Mary Whitfield,<sup>6</sup> Eben H. Paxton<sup>7</sup> and Thomas B. Smith<sup>1,8</sup>

### Abstract

Few regions have been more severely impacted by climate change in the USA than the Desert Southwest. Here, we use ecological genomics to assess the potential for adaptation to rising global temperatures in a widespread songbird, the willow flycatcher (*Empidonax traillii*), and find the endangered desert southwestern subspecies (*E. t. extimus*) most vulnerable to future climate change. Highly significant correlations between present abundance and estimates of genomic vulnerability – the mismatch between current and predicted future genotype–environment relationships – indicate small, fragmented populations of the southwestern willow flycatcher will have to adapt most to keep pace with climate change. Links between climate-associated genotypes and genes important to thermal tolerance in birds provide a potential mechanism for adaptation to temperature extremes. Our results demonstrate that the incorporation of genotype–environment relationships into landscape-scale models of climate vulnerability can facilitate more precise predictions of climate impacts and help guide conservation in threatened and endangered groups.

### Keywords

climate change, ecological genomics, genomic vulnerability, local adaptation.

Ecology Letters (2018) 21: 1085–1096

## INTRODUCTION

The effects of climate change on biodiversity are forecast to be one of the leading causes of extinction over the next century (Dawson *et al.* 2011; Warren *et al.* 2013; Pacifici *et al.* 2015; Urban 2015). Evidence of climate-induced local extinctions are now widespread among plant and animal species (Sinervo *et al.* 2010; Wiens 2016) and the velocity of climate change impacts in desert biomes is predicted to be among the fastest (Loarie *et al.* 2009). Recent climate change has altered community composition by favouring generalist taxa over habitat specialists and rare species (Menéndez *et al.* 2006; Estrada *et al.* 2016), but the ability to measure climate impacts below the species level is often lacking. Fine-scale estimates of vulnerability to climate change require an understanding of both the capacity for populations to shift their ranges to track climate conditions, as well as their capacity to tolerate climate alterations *in situ* via phenotypic plasticity or adaptation. Although intraspecific variation in climate tolerances may factor critically in the ability of species to move or adapt to environmental change, most modelling efforts ignore local adaptation. However, genomic tools are facilitating assessments of local adaptation in non-model species with increasing reliability (Savolainen *et al.* 2013) and such

information can be used to improve climate vulnerability estimates. Here, we combine genome-wide sequencing with environmental data to improve predictions of how genotype–environment relationships may be disrupted by future environmental change in an endangered songbird native to the Desert Southwest of the USA, the southwestern willow flycatcher.

Until recently, assessing species vulnerability to climate change focused largely on using current range–climate associations to predict distributions under models of future climate (Parmesan & Yohe 2003; Pacifici *et al.* 2015). However, complex biotic interactions (competition, specialisation, co-evolution, etc.) and or limits to dispersal imposed by physical barriers may limit range shifts, making it important to understand a species' potential to adapt to climate change *in situ* (Williams *et al.* 2008). Methodologies in the field of ecological genomics have provided tools to help incorporate information on local adaptation into climate vulnerability models by identifying regions where climate-induced selective pressure will be highest (Fitzpatrick & Keller 2015), but such methods have yet to be widely implemented. These approaches calculate the difference between current genotype–environment relationships and those predicted under future climate change to identify the geographic regions of greatest mismatch. More

<sup>1</sup>Center for Tropical Research, Institute for the Environment and Sustainability, University of California Los Angeles, Los Angeles, CA 90095, USA

<sup>2</sup>Department of Ecology and Evolutionary Biology, University of California Santa Cruz, Santa Cruz, CA 95060, USA

<sup>3</sup>Department of Evolution and Ecology, University of California Davis, One Shields Ave, Davis, CA 95616, USA

<sup>4</sup>Southwest Fisheries Science Center, National Marine Fisheries Service, 110 Shaffer Road, Santa Cruz, CA 95060, USA

<sup>5</sup>The Institute for Bird Populations, PO Box 1346, Point Reyes Station, CA 94956, USA

<sup>6</sup>Southern Sierra Research Station, P.O. Box 1316, Weldon, CA 932883, USA

<sup>7</sup>U.S. Geological Survey Pacific Island Ecosystems Research Center, Hawaii Volcano National Park, HI 96718

<sup>8</sup>Department of Ecology and Evolutionary Biology, University of California, Los Angeles, 621 Charles E. Young Drive South, Los Angeles, CA 90095, USA

\*Correspondence: E-mail: kruegg@ucsc.edu

†These authors contributed equally to this work.

specifically, they can be used to ask, ‘How much would allele frequencies across the range have to change to keep pace with projected changes in climate?’. In the absence of a range shift, populations in regions where the mismatch is greatest may either need to adapt or suffer population declines, as was recently shown in the North American songbird, the Yellow warbler (*Setophaga petechia*) (Bay *et al.* 2018).

Few regions in North America will be more severely impacted by temperature extremes than the Desert Southwest (Diffenbaugh *et al.* 2008; Hsiang *et al.* 2017). While most large-scale analyses of climate impacts in birds have focused on changes in geographic ranges or shifts in migratory phenology to better synchronise arrival times with earlier spring onset (Both & Visser 2001; Both *et al.* 2006; Stephens *et al.* 2016), these changes will do little to offset the impact of summer heat waves in desert regions. Recent work suggests that small desert passerines, in particular, will experience higher rates of mortality due to dehydration and hyperthermia as the frequency of extreme temperature events increases (Albright *et al.* 2017). In addition, work in poultry has shown that high temperatures can cause heart strain, or in some cases heart failure, as birds attempt to dissipate heat through increased blood circulation. Furthermore, this work has shown that such stress is not just physiological in nature, but is associated with differential expression in a suite of *c.* 300 genes (Zhang *et al.* 2017). Based on these studies, we predict that genes important to thermal cooling will be under strong selection in small desert passerines as the frequency of heat waves increases.

The endangered southwestern willow flycatcher provides an example of a desert passerine for which a better understanding of climate vulnerability has important implications for its conservation. This desert subspecies is one of four subspecies within the willow flycatcher whose combined ranges span the continental USA (Fig. 1; Pacific Northwestern form, *E. t. brewsteri*; Western Central form, *E. t. adastus*; and Eastern form, *E. t. trallii*). The presence of the southwestern willow flycatcher in particular is associated with riparian woodlands along streams and waterways (Sedgwick 2000) and such habitats are thought to provide important refuges from temperature extremes (Chen *et al.* 1999; McLeod *et al.* 2008). At the turn of the century, the southwestern willow flycatcher was described as common wherever its specialised habitat existed (Grinnell & Miller 1944), but by 1995 when it was listed under the Endangered Species Act, the number of known breeding pairs had been reduced to between 300 and 500 (Unitt 1987; Sogge *et al.* 1997). Population declines have been attributed to loss of riparian habitats in the Southwest following dam-building, water diversions, groundwater pumping, urbanisation, agricultural development and livestock grazing (Service 2002), but the role that climate change may have played in declines is unknown. Some researchers have questioned the subspecies designation of the southwestern willow flycatcher, suggesting that it is a peripheral population of an otherwise widespread species with no evidence for ecological distinctiveness (Zink 2015), although this suggestion has been questioned (Theimer *et al.* 2016). Here, we use ecological genomics to investigate the potential for ecological


distinctiveness within the willow flycatcher as well as the potential role of rising global temperatures on its future persistence.

To investigate potential genomic signals of local adaptation in the willow flycatcher, we tested for significant genotype–environment correlations using 105 000 SNP markers from 219 individuals spanning 24 populations across the breeding range (Fig. 1; Table 1). To identify the genomic locations of climate-associated SNPs in relation to genes and gene regions potentially important to adaptation under climate change, we also assembled and annotated the first willow flycatcher genome. Significant genotype–environment correlations for a subset of loci were further validated by genotyping an additional 274 individuals spanning mostly new 25 populations. To identify geographical regions where the mismatch between current and future genotype–environment relationships is predicted to be the greatest we used gradient forest modeling to calculate an index of genomic vulnerability (Bay *et al.* 2018). Lastly, we assessed the relationship between our estimates of genomic vulnerability and abundance across the range in order to assess which subspecies may be most vulnerable to future climate change.

## MATERIALS AND METHODS

### Sample collection and DNA extraction

We compiled a collection of 493 willow flycatcher blood or tissue samples from 41 locations across the breeding range using a combination of samples from previous studies, museum donations and new field collections (Paxton 2000).



**Figure 1** Willow Flycatcher Range Map and Sampling. Open and closed circles represent the data used in distance matrix comparison tests, while only populations represented by closed circles were used in the Gradient Forest analysis. Open grey boxes represent populations used to validate gene–environment correlations. Lines represent currently recognised subspecies boundaries according to Sogge *et al.* (1997). *E. t. brewsteri* = Pacific Coastal, *E. t. adastus* = Interior West, *E. t. trallii* = East and *E. t. extimus* = Southwest.

**Table 1** Sample location information.  $N_{\text{RAD\_nofilter}}$  = number of individuals for genome-wide RAD dataset before filtering for read depth and missing data,  $N_{\text{RAD\_filter}}$  = number of remaining post-filtering,  $N_{\text{validation}}$  = number of individuals in the SNP validation dataset

Location	Latitude	Longitude	$N_{\text{RAD\_nofilter}}$	$N_{\text{RAD\_filter}}$	$N_{\text{validation}}$
East Pima, AZ	32.83	-109.7	–	–	8
San Pedro/Gila River confluence, AZ	32.98	-110.77	18	14	17
West Fort Ditch, NM	33.04	-108.54	–	–	11
San Carlos Reservation, AZ	33.2	-110.44	–	–	30
San Diego, CA	33.28	-117.37	14	4	6
Roosevelt Lake, AZ	33.77	-111.24	20	10	18
White Mountains, AZ	34.00	-109.00	15	13	15
Camp Verde, AZ	34.56	-111.84	–	–	17
Santa Ynez River, CA	34.62	-120.18	–	–	8
Zuni/Nutria Diversion Reservation, NM	35.24	-108.64	–	–	8
South Fork Kern River, CA	35.66	-118.46	20	13	11
Southern Ute Reservation, CO	37.12	-107.59	–	–	6
Pahranagat Lake NWR, NV	37.32	-115.13	–	–	6
Owen's River at Bishop, CA	37.41	-118.48	–	–	12
Alamosa National Wildlife Refuge, CO	37.5	-106	–	–	17
Beaver Creek, CO	37.68	-108.38	–	–	6
Clear Creek, CO	37.79	-108.24	–	–	8
Baltimore Area, MD	39.4	-76.99	–	–	8
Escalante State Wildlife Area, CO	39.47	-106.37	–	–	13
Fish Creek, UT	39.78	-111.20	14	11	–
Rio Blanco Lake, CO	40.09	-108.21	–	–	7
Orefield, PA	40.66	-75.67	21	21	–
White River Confl. to the Green River, UT	40.67	-109.68	7	6	–
Willow Slew, IN	40.98	-87.53	4	4	–
Bigelow Meadows, CA	41.26	-121.88	7	6	–
Agusta, MI	42.3	-85.32	–	–	9
Mink Creek, ID	42.75	-112.39	6	6	–
Malheur NWR, OR	42.83	-118.87	7	6	–
FCTC-SABO, MI	42.84	-85.30	4	4	6
Jones Creek, OR	43.04	-123.97	10	10	–
Little White River Rec. Area, SD	43.17	-101.53	4	4	6
Black Creek, NY	43.38	-73.91	6	4	–
Fall Creek 2, ID	43.43	-111.40	7	7	–
Marion Forks, OR	44.37	-122.02	–	–	14
Finley NWR, OR	44.41	-123.35	3	0	–
Priem Road, OR	44.78	-123.38	7	6	–
Elm Creek, MN	45.13	-93.45	4	0	6
Waubay NWR, SD	45.40	-97.33	4	4	–
Hamon Memorial, MT	45.95	-114.13	5	4	–
Carbondale (Edgwick), WA	47.09	-122.05	8	7	–
Fork clearcut, WA	47.97	-124.40	4	4	–
Total			219	168	273

Two hundred and nineteen individuals from 24 populations were used to test for genome-wide genotype–environment correlations, while 274 individuals spanning 25 populations (eight replicate and 17 new populations) were used to validate a subset of significant genotype–environment correlations identified in the genome-wide analysis ( $N_{\text{total\_indiv}} = 493$ ;  $N_{\text{total\_pops}} = 41$ ; Table 1; Fig. 1). The willow flycatcher range map and associated subspecies boundaries was taken from the most current United States Geological Survey map used for willow flycatcher surveys (Sogge *et al.* 1997). Samples within one degree latitude and longitude and with no more than 10% difference

in any environmental variable (as indicated by our environmental analysis, below) were lumped into a single population. DNA was purified using the Qiagen™ DNeasy Blood and Tissue extraction kit and quantified using the Qubit® dsDNA HS Assay kit (Thermo Fisher Scientific, USA).

#### Genome sequencing, assembly and annotation

The genomic DNA library was created using a single southwestern willow flycatcher individual from Roosevelt Lake, AZ and the Illumina TruSeq DNA PCR-Free LT kit (Illumina,

Hayward, CA), with adjustments. One  $\mu\text{g}$  of DNA was diluted in 100  $\mu\text{L}$  of AE buffer and fragmented to an average insert size of *c.* 400 bp. The resulting library was sequenced on two lanes of an Illumina HiSeq2500 using 250 bp paired-end sequencing at the QB3 Vincent J. Coates Genomics Sequencing Laboratory, UC Berkeley. Two mate-pair libraries were also created, using 4 and 8 kb inserts and sequenced on one-third of a Illumina HiSeq 2500 lane, using 100 bp paired-end sequencing at the Huntsman Cancer Center at the University of Utah. The 250 bp paired-end reads were used to assemble contigs with the Discovar DeNovo assembler from the Broad Institute (<http://www.broadinstitute.org>), discarding contigs less than 1000 bp in length. Mate-pair reads were trimmed and separated from paired-end reads using NxTrim (O'Connell *et al.* 2015) and contigs were scaffolded with SSPACE (overlap requirement  $k = 3$ ) (Boetzer *et al.* 2010) using both paired-end and mate-pair libraries. We used reapr (Hunt *et al.* 2013) and mapping of the 8 kb insert library to break the assembly at likely error regions. SSPACE scaffolding was repeated with  $k = 5$  and scaffolds  $< 5\text{ kbp}$  were discarded for the final assembly.

For annotation purposes, repetitive regions were replaced with N's using RepeatMasker (-species birds) (Tarailo-Graovac & Chen 2009). For annotation, we used two different *ab initio* gene predictions within the MAKER pipeline (Cantarel *et al.* 2008): SNAP and AUGUSTUS. SNAP was trained iteratively using Zebra Finch cDNA and protein sequences downloaded from Ensembl and AUGUSTUS was run using the available chicken training dataset. We used Interproscan (Zdobnov & Apweiler 2001) to add Pfam protein annotation and gene ontology (GO) terms and identified 15 489 genes. Scaffolds were aligned to the Zebra Finch genome (version 3.2.4) using the software promer, part of the MUMmer package (Delcher *et al.* 2003). After alignment, we retained the longest consistent alignment (-q) for each chromosome while filtering for similarity (-i 50) and alignment length (-l 500). We then determined the location of the longest alignment for each scaffold and ordered scaffolds accordingly for visualisation purposes.

### SNP discovery and SNP filtering

Genome scans on 219 individuals were conducted following the BestRAD library preparation protocol with some modifications (Ali *et al.* 2016). 100 ng of DNA was digested using the SbfI restriction enzyme (New England Biolabs, USA) and fragments were ligated with SbfI adapters prepared with biotinylated ends. Adapter-ligated samples were pooled and cleaned using 1X Agencourt<sup>®</sup> AMPure XP beads (Beckman Coulter Inc., USA). All DNA fragments were sheared to an average length of 400 bp and adapter-ligated fragments were bound to M-280 streptavidin magnetic Dynabeads (Life Technologies, USA). Blunt end repair and ligation of NEBNext Adapters was performed using the Illumina NEBNext Ultra DNA Library Prep Kit (New England Biolabs, USA) and Agencourt<sup>®</sup> AMPure XP beads (Beckman Coulter Inc, USA) were used to size select an average of 500 bp fragments. The final library was cleaned and run on a Bioanalyzer at the UCLA Technology Center for Genomics & Bioinformatics to check for the size distribution and the absence of

contaminants. Two libraries, each comprised of 96 individuals, were initially sequenced in two lanes of 100 bp paired-end reads on an Illumina HiSeq 2500 at the UC Davis Genome Center. In a third lane, 69 individuals with low coverage from the first two libraries were re-sequenced along with an additional 27 new individuals.

The program Stacks (Catchen *et al.* 2013) was used to demultiplex, filter and trim adapters from the data with the process\_radtags function and to remove duplicate read pairs using the clone\_filter function. Reads were mapped to our genome assembly using bowtie2 (Langmead & Salzberg 2012) and the Haplotype Caller in the Genome Analysis Toolkit was used to identify single nucleotide polymorphisms (SNPs), following best practices from the Broad Institute (<http://www.broadinstitute.org>). Finally, we discarded low-quality and rare variants (genotype quality  $< 30$ ; depth  $< 8$ ; minor allele frequency  $< 0.01$ ), as well as indels and non-biallelic SNPs using vcftools (Danecek *et al.* 2011). We used the R package genoscopeRtools (<https://doi.org/10.5281/zenodo.848279>) to visualise the tradeoff between discarding SNPs with low coverage and discarding individuals with missing genotypes in order to determine the final number of SNPs and individuals retained (Fig. S1).

### Environmental data

For each sampling location, we obtained environmental data from publicly available databases. These 25 variables included 19 climate variables downloaded from WorldClim (Hijmans *et al.* 2005) which represented average climate between the years 1960 and 1990, as well as vegetation indices (Carroll *et al.* 2004) (NDVI and NDVIstd, average for the year 2003), Tree Cover (Sexton *et al.* 2013) and elevation data from the Global Land Cover Facility (<http://www.landcover.org>) and a measure of surface moisture characteristics from the NASA Scatterometer Climate Record Pathfinder project (QuickSCAT mean and standard deviation, downloaded from [scp.byu.edu](http://scp.byu.edu)).

### Assessing the role of geography and environment

To assess the relative contributions of geography and the environment to genetic divergence in the willow flycatcher, we compared genetic, environmental and geographic distance matrices and used multiple tests designed to account for spatial autocorrelation. For locations with  $> 4$  individuals (Table 1), we calculated pairwise  $F_{ST}$  across all quality-filtered SNPs using the R package SNPrelate (Zheng *et al.* 2012) and pairwise geographic distances from longitude and latitude using the R package geosphere (Hijmans *et al.* 2012). We then calculated environmental distance between each pair of sites by removing highly correlated climate variables (Pearson's  $r > 0.7$ ; Table 2; Table S1), scaling and centring each environmental variable to account for differences in magnitude, and then calculating pairwise Euclidean differences between sites. Mantel, Partial Mantel and multiple regression of distance matrices were used to test for associations between linearised  $F_{ST}$  ( $F_{ST}/1-F_{ST}$ ) and genetic and environmental distance after accounting for geographic distance.



**Table 2** Environmental variables used in the gradient forest analysis, ordered by ranked importance of variables and the cumulative contribution of each variable. The top eight environmental variables represent 49% of the total

Variable	Definition	GF rank	Cumulative contribution
BIO11*	Mean temperature of coldest quarter	8.03E-04	7.66
BIO10	Mean temperature of warmest quarter	6.71E-04	14.40
BIO1	Annual mean temperature	6.41E-04	21.05
BIO5*	Max temperature of warmest month	6.40E-04	27.47
BIO6	Min temperature of coldest month	5.79E-04	32.90
BIO4*	Temperature seasonality (standard deviation *100)	5.20E-04	38.30
BIO9	Mean temperature of driest quarter	4.91E-04	43.64
BIO17*	Precipitation of driest quarter	4.78E-04	48.76
NDVI_Mean	Vegetation indices	4.50E-04	53.18
BIO15	Precipitation seasonality (Coefficient of Variation)	4.28E-04	57.39
BIO7	Temperature annual range (BIO5-BIO6)	3.75E-04	61.41
TreeCover	Tree cover	3.72E-04	65.41
BIO14	Precipitation of driest month	3.64E-04	69.36
BIO16	Precipitation of wettest quarter	3.09E-04	72.75
BIO19	Precipitation of coldest quarter	2.96E-04	76.04
BIO2*	Mean diurnal range (Mean of monthly (max temp – min temp))	2.90E-04	79.31
BIO8*	Mean temperature of wettest quarter	2.87E-04	82.49
BIO13	Precipitation of wettest month	2.82E-04	85.63
STM	Elevation	2.21E-04	88.36
BIO12	Annual precipitation	2.12E-04	90.98
BIO3	Isothermality (BIO2/BIO7) (* 100)	2.07E-04	93.54
QuickScat	Surface moisture characteristics	2.02E-04	95.87
BIO18	Precipitation of warmest quarter	1.92E-04	98.17
NDVI_StDev	Vegetation indices	1.81E-04	100.00

\*Top-ranked, uncorrelated climate variables used for Gradient Forest mapping and distance matrix comparison analyses.

These variables were selected by moving down the list of ranked importance for the full model and discarding variables highly correlated (Pearson's  $r > 7$ ) with a variable of higher importance.

### Gradient forest prediction of genomic mismatch

We identified the environmental variables that best explained genetic variation using gradient forest analysis with the *R* package gradientForest (Ellis *et al.* 2012). Because rare alleles are more likely to yield false positives, we only used SNPs with minor allele frequency > 10%. The gradient forest analysis (nree = 500, nbin = 201, corr.threshold = 0.5) provided a ranked list based on the relative predictive power of all environmental variables (Table 2). To ensure that our model was explaining more variation than we would expect by chance, we compared the number of SNPs with positive  $R^2$  and the mean  $R^2$  across these 'predictive' loci (those with positive  $R^2$ ) to 10 runs with randomised environments. Visualisation of the gradient forest model across the range of the willow flycatcher (Buschke *et al.* 2016) was done by generating and extracting uncorrelated BIOCLIM values for 100 000 random points. The final gradient forest model was used to predict the genomic composition from uncorrelated environmental variables for each random point (Table 2). Principal components analysis (PCA) was used to summarise values. To visualise the different adaptive environments across the breeding range, colours were assigned based on the top three principal components axes, as recommended by the authors (Ellis *et al.* 2012).

We extended the gradient forest analysis to predict 'genomic vulnerability' using the method presented by Fitzpatrick & Keller (2015). Here, 'genomic vulnerability' (termed 'genetic offset' by Fitzpatrick and Keller) is a measure of the mismatch between genotype and future predicted environment

using associations across current gradients as a baseline. We used the baseline gradient forest model calculated using current BIOCLIM values to predict genomes under future environmental conditions (based on RCP 2.6 2050 projections) at the same 100 000 random points. The Euclidean distance between these weighted current and predicted values is what we refer to as 'genomic vulnerability' (Bay *et al.* 2018).

### Identification of SNPs as candidates for environmental selection

To identify SNPs (with minor allele frequency > 0.1) that were most highly associated with the top environmental variables while accounting for underlying population structure, we used Latent Factor Mixed Models (LFMM) (Frichot *et al.* 2013). For each of the top eight environmental variables from the gradient forest analysis, we ran five separate MCMC runs with a latent factor of  $K = 4$ , based on the number of reported subspecies and previous morphological and genetic analysis based upon neutral markers (Paxton 2000). *P*-values from all five runs were combined and adjusted for multiple tests using a false discovery rate (FDR) correction. We annotated each significant SNP with genes within 25 kb upstream or downstream which we assume is within the distances before which LD should break down (Backstrom *et al.* 2006).

### Validation of climate-associated SNPs

To validate genotype–environment correlations identified in the LFMM analysis, we genotyped the top-ranking 18 SNPs

that were significantly associated with the top eight climate variables and could be converted to SNPtype Assays in an additional 274 breeding individuals from 25 locations. DNA was extracted from feather samples using the KingFisher™ Cell and Tissue DNA Kit and SNP genotyping was performed on the Fluidigm™ (ThermoFisher Scientific, USA) 96.96 IFC controller (Fluidigm Inc., San Francisco, CA, USA) following manufacturer guidelines. Nine individuals with > 8% of missing data were removed from downstream analysis and final allele frequencies were calculated for each SNP at each location. Standard linear regression was used to test for significant associations between climate and allele frequency (FDR-corrected  $P$ -value < 0.05).

#### Association between genomic vulnerability and abundance

To assess the relationship between genomic vulnerability and abundance and determine which subspecies may be most vulnerable to future climate change, we correlated estimates of genomic vulnerability with willow flycatcher relative abundance from the North American Breeding Bird Survey (BBS) for 2011–2015, including all sites where the species was detected at least once during the history of the survey (Pardieck *et al.* 2017). In order to associate the two datasets, vector-based BBS relative abundance estimates derived from inverse-distance weighting interpolation (2010–15; Sauer *et al.* 2017; [https://www.mbr-pwrc.usgs.gov/bbs/shape\\_ra15.html](https://www.mbr-pwrc.usgs.gov/bbs/shape_ra15.html)) of route-level mean counts were converted to raster format with grid resolution of approximately  $15 \times 15$  km. We then extracted values of relative abundance and genomic vulnerability for grid cells including BBS routes using bilinear interpolation (Hijmans 2015). For cells with BBS routes where detections had been recorded, but for which model-based estimates of abundance were not available due to low abundance and isolation from other sites with detections, we assigned mean count values (*c.* 9% of routes; mean count = 0.06). Significant differences in genomic vulnerability between subspecies were assessed using boxplots with 95% confidence intervals around median vulnerability scores (Chambers *et al.* 1983).

## RESULTS

#### Genome assembly, SNP discovery and SNP/population filtering

The final southwestern willow flycatcher genome assembly was 1.2 Gb in length and consisted of 7791 scaffolds (contig N50 = 79 613 bp; scaffold N50 = 895,074 bp). In total, we identified 6 355 061 SNPs across the genome. Discarding low quality SNPs and low-coverage individuals resulted in a final set of 105 000 SNPs and 175 individuals (Fig. S1), with less than 7.4% missing genotypes per SNP (mean = 2.3%), < 15.6% missing SNPs per individual (mean = 2.3%) and minor allele frequency greater than 1%. Because  $F_{ST}$  is robust to low sample size when a large number of SNPs are employed (Nazareno *et al.* 2017), we retained all populations with a minimum of four (mean = 8) individuals for analysis based upon  $F_{ST}$  (distance matrix comparisons), resulting in a final dataset of 168 individuals from 22 sampling locations.

Alternatively, to avoid bias associated with low sample size in analyses requiring estimates of allele frequency (Gradient Forest and LFMM), we used only populations with a minimum of six individuals (average = 10), resulting in a final dataset of 136 individuals from 14 sampling locations (Fig. 1; Table 1).

#### Assessing the role of geography and environment in shaping genetic structure


Pairwise  $F_{ST}$  across all quality-filtered SNPs ranged from 0 to 0.11 (Table S2). Mantel tests revealed highly significant correlations between genetic and geographic distance ( $r = 0.70$ ,  $P = 1 \times 10^{-6}$ ), genetic and environmental distance ( $r = 0.56$ ,  $P = 1 \times 10^{-6}$ ) and geographic and environmental distance ( $r = 0.42$ ,  $P = 1.8 \times 10^{-4}$ ) (Fig. S2A). Partial Mantel tests revealed the correlation between genetic and environmental distance remained significant after accounting for the relationship between genetic and geographic distance ( $r = 0.42$ ,  $P = 3 \times 10^{-4}$ ; Fig. S2), and both geographic and environmental distances were significant in a multiple regression of distance matrices (MRM:  $R^2=0.59$ ; geography  $P = 1 \times 10^{-5}$ ; environment  $P = 3 \times 10^{-5}$ ).

#### Gradient forest mapping of genotype–environment correlations

More genetic variation was explained by our gradient forest than those generated under randomised environments (Fig. S3). A total of 9015 SNPs were correlated with environment with mean  $R^2 = 0.18$ , compared to a mean  $R^2$  of 0.13–0.15 across 3489–5633 SNPs for randomised data. We used gradient forest models to identify which climate and vegetation variables were most important in structuring genetic variation in the willow flycatcher and visualise climate-associated allelic variation across the breeding range (Fig. 2a and b). Seven temperature variables and one precipitation variable were most strongly correlated with genetic variation across the breeding range of willow flycatchers (Table 2). Mapping principal components of gradient forest output revealed putative signals of local adaptation across the US Southwest, the East, the Inter-Mountain West and the Pacific Northwest geographic regions (Fig. 2c).

#### Identification of candidate SNPs for environmental selection

To investigate genomic regions potentially involved in climate adaptation, we identified genomic regions associated with the top eight climatic variables (which explained 49% of the total variation) using Latent Factor Mixed Models (25) (Table 2, Table S3). We found 77, 100, 104, 97, 97, 58, 107 and 70 SNPs significantly associated with BIO11, BIO10, BIO5, BIO1, BIO6, BIO9, BIO4 and BIO17, respectively (FDR-corrected  $P < 0.05$ ), with one SNP located on chromosome 16, Climate\_20, shared among seven variables. The SNPs were broadly distributed across the genome and within 25 KB of 202 genes with a variety of functions (Table S3). We identified five genes (BRACA1, RND2, CIITA, ICOS and UBE2C) that were among the *c.* 300 genes found to be differentially expressed in an RNA-seq analysis of thermal tolerance in chickens (Zhang *et al.* 2017), two of which were physically



**Figure 2** Mapping gene–environment correlations across the willow flycatcher breeding range. (a) Principal components analysis of gradient forest-transformed climate variables. Black dots represent the PC scores associated with the sampling locations, while colours are based upon modelled gene–environment correlations from 100 000 random points across the breeding range. Arrows show the loadings of the top-ranked uncorrelated environmental variables. (b) Gradient forest-transformed climate variables from the PCA mapped to geography support climate adaptation across the breeding range. Black lines designating approximate subspecies locations support the idea that while subspecies are adapted to distinct ecological regions, climate adaption is complex.

linked (BRACA1 and RND2), and an additional five genes (Ecel1, SLC23A2, NOX4, PIRT and GR1N1) with GO terms related to other aspects of thermal tolerance, including respiratory system process, oxidative stress and response to heat (Rimoldi *et al.* 2015) (Table S4). Three of the five genes from the poultry thermal stress study were found to be outliers in association with BIO4, Temperature Seasonality (Fig. 3a). Furthermore, targeted genotyping using Fluidigm SNPtype assays for 18 of the top candidate SNPs in an additional 274 birds from 24 locations validated climate associations in 8/18 SNPs (FDR-corrected  $P < 0.05$ ; Table S5). In particular, we found a highly significant relationship between the Climate\_20 SNP and seven of the eight top-ranked climate variables in both the genome scan and validation results. While no link between Climate\_20 and genes linked to thermal tolerance in birds was found, the highly significant relationship between this SNP and climate variables reflective of the intensity of summer heat waves, such as Mean Temperature of the Warmest Quarter (BIO10), suggests a potential role for this region in climate adaptation (Fig. 3b, c and d).

#### Prediction of genomic mismatch and association between vulnerability and abundance


Under a model of future climate change, genomic vulnerability was predicted to be highest in the southern part of the willow flycatcher range (Fig. 4a), corresponding to the range of the southwestern willow flycatcher subspecies range. Overall, highest genomic vulnerability occurred at sites with especially low abundance, resulting in a significant negative correlation between abundance and genetic vulnerability ( $r = -0.18$ ;  $P < 0.001$ ; d.f. = 1382; Fig. 4b, c). Abundance of southwestern willow flycatcher was low across sites and correlation

between abundance and vulnerability for this subspecies was especially strong ( $r = -0.49$ ;  $P = 0.016$ ; d.f. = 27) and weakest for the eastern subspecies region (*trillii*;  $r = -0.11$ ;  $P < 0.001$ ; d.f. = 957). While there were regions of high and low genomic vulnerability across the range, the southwestern willow flycatcher subspecies had the highest overall median genomic vulnerability score (Fig. 4d).

#### DISCUSSION

Climate envelope models are widely used to predict future species distributions (Parmesan & Yohe 2003; Pacifici *et al.* 2015), but such models do not account for complex biotic interactions (competition, specialisation, co-evolution, etc.) or barriers to dispersal that may limit range shifts (Williams *et al.* 2008). In the case of the willow flycatcher, the capacity for range shifts may be restricted by the need to be proximate to specific water sources (Figgins & Finch 2015), making it important to incorporate the potential for adaptation into estimates of climate vulnerability. Here, we move beyond species distribution modelling to begin to identify populations that will need to adapt most to keep pace with climate change. By calculating the difference between current genotype–environment relationships and those predicted under future climate change, we identify regions of highest vulnerability in the southern part of the range. A comparison of the average genomic vulnerability across all currently recognised subspecies strongly supports the view that the endangered southwestern willow flycatcher is most vulnerable to climate change. Significant correlations between estimates of genomic vulnerability and abundance from Breeding Bird Survey data confirm that already rare populations in the Southwest and throughout the range have the highest genomic vulnerability, suggesting that climate






**Figure 3** Candidate SNPs linked to temperature in the Willow flycatcher. (a) Manhattan plot showing the FDR-corrected significance level for SNPs associated with Temperature Seasonality (BIO4) and (b) Mean Temperature of the Warmest Quarter (BIO10). Dashed line represents  $P = 0.05$ . Colours distinguish different chromosomes. Candidate genes linked to thermal tolerance in birds are highlighted by red stars and denoted with gene names, while Climate\_20, the SNP validated in B and C below, is denoted by a black triangle. No link between Climate\_20 and genes linked to thermal tolerance in birds was found, but the highly significant relationship between this SNP and seven of the eight top-ranked climate variables (except temperature seasonality shown in A above) in both the genome scan and validation results (Table S5) suggest a potential role for this region in climate adaptation. (c) Relationship between Climate\_20 and mean temperature of the warmest quarter in genome scan and SNP validation datasets. The allele frequencies from the original genome scan data are denoted by squares, while allele frequencies based upon the validation set are denoted by circles. (d) The association between Mean Temperature of the Warmest Quarter (BIO10) and Climate\_20 across geographic space, with population allele frequencies colour coded from high frequency (red) to low (yellow).

change may have already had an impact on population declines in regions at the edge of the species niche. Our results demonstrate how the incorporation of genotype–environment relationships into models of climate vulnerability can improve predictions of climate-induced impacts below the species level.

Assessing the extent of intraspecific variation in climate tolerances is an important first step towards understanding species vulnerability to climate change. Here, we investigate the relationship between genetic, geographic and environmental distance in the willow flycatcher and find consistent support for the conclusion both geography and environment are

important to genetic divergence in the willow flycatcher (Fig. S2). Mapping putatively adaptive genetic variation using gradient forest-transformed climate variables supports the idea that the Pacific Northwest, the Southwest, the East, and the Inter-Mountain West harbour unique genotype–environment correlations. More specifically, our results support the idea that high maximum temperatures during the warmest month (BIO5) are important to genotype–environment correlations in the Southwest, while genotype–environment relationships in the Pacific Northwest are driven by environmental variables such as precipitation during the driest quarter (BIO17) and mean temperatures during the coldest quarter (Figs 2 and 3).





**Figure 4** Genomic Vulnerability and abundance in the Willow Flycatcher. (a) Map of genomic vulnerability across the Willow Flycatcher breeding range. Red = high genomic vulnerability, blue = low genomic vulnerability, lines indicate subspecies boundaries. (b) Genomic Vulnerability vs. abundance based upon the estimated mean number of birds/ route in 2011–2015 Breeding Bird Survey. (c) Estimates of relative abundance from the BBS based on inverse-distance weighting interpolation. Points indicate the BBS routes where Willow Flycatchers have been recorded. Points in the grey areas fall in regions where abundance was too low or distant from other detection routes to be included in the BBS spatial model. (d) Quantile box plots of the median Genomic Vulnerability broken down by subspecies. Open circles represent outliers.

In contrast, genotype–environment correlations in Inter-Mountain West and Eastern populations, centre closer zero in the PCA (Fig. 2a), indicating a more moderate impact of climate variables underlying climate adaptation in this area. In sum, our results support the idea that genotype–environment correlations in the willow flycatcher are complex, involving multiple environmental variables and genomic regions and such information can be used to help refine estimates of future climate vulnerability.

Adaptation to local environments often occurs through natural selection acting on a large number of loci, each with a small effect on phenotype (Orr 2005). Here, we identify putative loci important to local adaptation in the willow flycatcher, after accounting for underlying population structure, and find between 58 and 107 SNPs significantly associated with each of the top eight environmental variables (Table S3). Independent validation of our top climate-associated SNPs in 274 new individuals from 24 populations revealed that eight of our top 18 loci were likely robust to Type 1 error. While such error is a problem common to all association studies (McCarthy *et al.* 2008), the high number of false positives in our data underscore the idea that genotype–environment associations that cannot be validated should be interpreted with caution. Highly significant associations between Climate<sub>20</sub> and seven of our eight top-ranked environmental variables in both the genome scan and validation datasets provide the strongest evidence for local adaptation across the willow

flycatcher genome (Fig. 3). While no associations between Climate<sub>20</sub> and genes known to be important to thermal tolerance in birds were identified, the relationship between allele frequency variation in this SNP and Mean Temperature of the Warmest Quarter (BIO5) suggests a potential role for this region in adaptation to temperature extremes. Overall, our results are in keeping with the idea that willow flycatchers exhibit region-specific genotype–climate associations that should be considered when assessing the capacity for endangered populations of the southwestern willow flycatcher to shift their range in response to rising global temperatures.

While genotype–environment correlations have been noted across a variety of plant and animal systems, the mechanisms behind such local adaptation remain less well understood. Recent work on birds supports the idea that exposure to high temperatures can result in dehydration and heat stress related mortality (Albright *et al.* 2017; Zhang *et al.* 2017). As a first step towards understanding the genomic basis of adaptation to temperature in the willow flycatcher, we identify genes within 25KB of our top ranking climate-associated SNPs (Table S4). Our strongest evidence for genes and gene regions that may be important to climate adaptation in this species comes from the overlap between five genes in our panel (BRACA1, RND2, CIITA, ICOS and UBE2C) and those that were also found to be differentially expressed in a thermal tolerance study in poultry (Zhang *et al.* 2017). More specifically, Zhang *et al.* (2016) concluded that expression of these genes

was linked to the dissipation of heat through increased heart pumping and blood circulation in smaller breeds of chickens. These results are consistent with the recent work by Albright *et al.* (2017) who found that small passerines in the Desert Southwest were particularly prone to mortality resulting from the failure to maintain body temperatures below lethal limits. While more research is needed, it is possible that physiological pathways responsible for overheating are related to those involved in interspecific adaptation to temperature extremes. Furthermore, while limited gene annotation information for non-model organisms makes us cautious about placing significance on GO term analyses (Stein 2001), we also note the presence of five genes (*Ecell1*, *SLC23A2*, *NOX4*, *PIRT* and *GRIN1*) with GO terms related to heat stress, thermal tolerance and oxidative stress. Future efforts will focus on validating gene-environment correlations at putative heat stress-related loci as well as investigating the extent to which the genes identified here may serve as a mechanism for adaptation to temperature extremes in the willow flycatcher.

Desert ecosystems are home to some of the world's rarest species, many of which are already threatened by climate change (Loarie *et al.* 2009). Methods for assessing climate change impacts that rely on single species distribution models may overlook the importance of local adaptation in the ability of populations to respond to environmental shifts, potentially leading to misplaced conservation efforts. The US Fish and Wildlife Service was considering removing the southwestern willow flycatcher from the endangered species list, in part because of a single species distribution model that showed no evidence of habitat specialisation across the range. Here, we annotate the first willow flycatcher genome and use population-level, genome-wide sequencing to show that willow flycatchers are not a single homogenous group, but a composite of locally adapted populations with specific genotype-environment relationships related to differences in temperature extremes. Clear evidence for local adaptation across the range highlights the need for management efforts below the species level if locally adapted populations are to be conserved. Estimates of the mismatch between current genotype-environment correlations and those predicted under future climate indicate that the southwestern subspecies is at the greatest risk of climate-induced extinction. Our findings support the idea that protection or enhancement of riparian thermal refuges (Chen *et al.* 1999) within regions of lower genomic vulnerability in the Desert Southwest may be the most effective strategy for conserving remaining populations of flycatchers by buffering them from temperature extremes.

#### ACKNOWLEDGEMENTS

We thank the many individuals who contributed genetic samples, including T. Kita, B. Kus, R. Taylor, M. Fylling and many MAPs (Monitoring Avian Productivity and Survivorship) station operators within the Institute for Bird Populations Network. This work used the Extreme Science and Engineering Discovery Environment (XSEDE), which is supported by National Science Foundation grant ACI-1548562. We thank the Vincent J. Coates Genomics Sequencing Laboratory at the University of California, Berkeley as well as the

UC Davis Genome Center for their help with the sequencing. This work was made possible by a generous gift from J. Ellis as well as an NSF Postdoctoral Fellowship (to R. Bay), a California Energy Commission grant EPC-15-043 (to K. Ruegg and T. Smith), a National Geographic grant WW-202R-17 (to K. Ruegg), and donation from First Solar Incorporated. Any use of trade, product, or firm names in this publication does not imply endorsement by the U.S. Government.

#### AUTHOR CONTRIBUTIONS

K.R., R.A.B. and T.B.S. conceived the study; R.A.B. assembled and annotated the genome; R.B., K.R., E.C.A., J.F.S. and R.J.H. contributed to the population genetic, BBS and landscape genetic analyses; M.W. and E.H.P. contributed samples and biological expertise; K.R. wrote the paper with contribution from all authors.

#### DATA ACCESSIBILITY STATEMENT

The Willow flycatcher genome and annotations are available through NCBI (accession number: PWAB00000000) and population-level RAD-Seq data are available through NCBI's Sequence Read Archive (<http://www.ncbi.nlm.nih.gov/bioproject/453612>).

#### REFERENCES

- Albright, T.P., Mutiibwa, D., Gerson, A.R., Smith, E.K., Talbot, W.A., O'Neill, J.J. *et al.* (2017). Mapping evaporative water loss in desert passerines reveals an expanding threat of lethal dehydration. *Proc. Natl Acad. Sci.*, 114, 2283–2288.
- Ali, O.A., O'Rourke, S.M., Amish, S.J., Meek, M.H., Luikart, G., Jeffres, C. *et al.* (2016). RAD capture (Rapture): flexible and efficient sequence-based genotyping. *Genetics*, 202, 389–400.
- Backstrom, N., Ovarnstrom, A., Gustafsson, L. & Ellegren, H. (2006). Levels of linkage disequilibrium in a wild bird population. *Biol Letters*, 2, 435–438.
- Bay, R., Harrigan, R.J., Underwood, V.L., Gibbs, H.L., Smith, T.B. & Ruegg, K. (2018). Genomic signals of selection predict climate-driven population declines. *Science*, 359, 83–89.
- Boetzer, M., Henkel, C.V., Jansen, H.J., Butler, D. & Pirovano, W. (2010). Scaffolding pre-assembled contigs using SSPACE. *Bioinformatics*, 27, 578–579.
- Both, C. & Visser, M.E. (2001). Adjustment to climate change is constrained by arrival date in a long-distance migrant bird. *Nature*, 411, 296–298.
- Both, C., Bouwhuis, S., Lessells, C. & Visser, M.E. (2006). Climate change and population declines in a long-distance migratory bird. *Nature*, 441, 81–83.
- Buschke, F.T., Brendonck, L. & Vanschoenwinkel, B. (2016). Adding energy gradients and long-distance dispersal to a neutral model improves predictions of Madagascar bird diversity. *Ecol. Evol.*, 6, 6919–6929.
- Cantarel, B.L., Korf, I., Robb, S.M., Parra, G., Ross, E., Moore, B. *et al.* (2008). MAKER: an easy-to-use annotation pipeline designed for emerging model organism genomes. *Genome Res.*, 18, 188–196.
- Carroll, M., DiMiceli, C., Sohlberg, R. & Townshend, J. (2004). 250 m MODIS normalized difference vegetation index. University of Maryland, College Park, Maryland.
- Catchen, J., Hohenlohe, P.A., Bassham, S., Amores, A. & Cresko, W.A. (2013). Stacks: an analysis tool set for population genomics. *Mol. Ecol.*, 22, 3124–3140.

- Chambers, J.M., Cleveland, W.S., Kleiner, B. & Tukey, P.A. (1983). *Graphical Methods for Data Analysis*. Wadsworth Belmont, CA.
- Chen, J., Saunders, S.C., Crow, T.R., Naiman, R.J., Broszofski, K.D., Mroz, G.D. *et al.* (1999). Microclimate in forest ecosystem and landscape ecology: variations in local climate can be used to monitor and compare the effects of different management regimes. *Bioscience*, 49, 288–297.
- Danecek, P., Auton, A., Abecasis, G., Albers, C.A., Banks, E., DePristo, M.A. *et al.* (2011). The variant call format and VCFtools. *Bioinformatics*, 27, 2156–2158.
- Dawson, T.P., Jackson, S.T., House, J.I., Prentice, I.C. & Mace, G.M. (2011). Beyond predictions: biodiversity conservation in a changing climate. *Science*, 332, 53–58.
- Delcher, A.L., Salzberg, S.L. & Phillippy, A.M. (2003). Using MUMmer to identify similar regions in large sequence sets. *Current Protocols in Bioinformatics*, 10, 10–3.
- Diffenbaugh, N.S., Giorgi, F. & Pal, J.S. (2008). Climate change hotspots in the United States. *Geophys. Res. Lett.*, 42, 3521–3528.
- Ellis, N., Smith, S.J. & Pitcher, C.R. (2012). Gradient forests: calculating importance gradients on physical predictors. *Ecology*, 93, 156–168.
- Estrada, A., Morales-Castilla, I., Caplat, P. & Early, R. (2016). Usefulness of species traits in predicting range shifts. *Trends Ecol. Evol.*, 31, 190–203.
- Fitzpatrick, M.C. & Keller, S.R. (2015). Ecological genomics meets community-level modelling of biodiversity: mapping the genomic landscape of current and future environmental adaptation. *Ecol. Lett.*, 18, 1–16.
- Frichot, E., Schoville, S.D., Bouchard, G. & François, O. (2013). Testing for associations between loci and environmental gradients using latent factor mixed models. *Mol. Biol. Evol.*, 30, 1687–1699.
- Friggens, M.M. & Finch, D.M. (2015). Implications of climate change for bird conservation in the southwestern US UNDER three alternative futures. *PLoS ONE*, 10, E0144089.
- Grinnell, J. & Miller, A.H. (1944). *The Distribution of the Birds of California*. The Cooper Ornithological Society, Berkeley, CA.
- Hijmans, R.J. (2015). Geographic Data Analysis and Modeling. R package version 2.5-8.
- Hijmans, R.J., Cameron, S.E., L., P.J., Jones, P.G. & Jarvis, A. (2005). Very high resolution interpolated global terrestrial climate surfaces. *Int. J. Climatol.*, 25, 1965–1978.
- Hijmans, R.J., Williams, E. & Vennes, C. (2012). Geosphere: Spherical Trigonometry. R package version 1.2–28. CRAN. R-project.org/package=geosphere.
- Hsiang, S., Kopp, R., Jina, A., Rising, J., Delgado, M., Mohan, S. *et al.* (2017). Estimating economic damage from climate change in the United States. *Science*, 356, 1362–1369.
- Hunt, M., Kikuchi, T., Sanders, M., Newbold, C., Berriman, M. & Otto, T.D. (2013). REAPR: a universal tool for genome assembly evaluation. *Genome Biol.*, 14, R47.
- Langmead, B. & Salzberg, S.L. (2012). Fast gapped-read alignment with Bowtie 2. *Nat. Meth.*, 9, 357–359.
- Loarie, S.R., Duffy, P.B., Hamilton, H., Asner, G.P., Field, C.B. & Ackerly, D.D. (2009). The velocity of climate change. *Nature*, 462, 1052.
- McCarthy, M.I., Abecasis, G.R., Cardon, L.R., Goldstein, D.B., Little, J., Ioannidis, J.P. *et al.* (2008). Genome-wide association studies for complex traits: consensus, uncertainty and challenges. *Nat. Rev. Genet.*, 9, 356.
- McLeod, M.A., Koronkiewicz, T.J., Brown, B.T., Langeberg, W.J. & Carothers, S.W. (2008). Southwestern Willow Flycatcher surveys, demography, and ecology along the lower Colorado River and tributaries, 2003–2007. Five-year summary report submitted to U.S. Bureau of Reclamation, Boulder City, Nevada. SWCA Environmental Consultants, Flagstaff, Arizona.
- Menéndez, R., Megías, A.G., Hill, J.K., Braschler, B., Willis, S.G., Collingham, Y. *et al.* (2006). Species richness changes lag behind climate change. *Proc. Biol. Sci.*, 273, 1465–1470.
- Nazareno, A.G., Bemmels, J.B., Dick, C.W. & Lohmann, L.G. (2017). Minimum sample sizes for population genomics: an empirical study from an Amazonian plant species. *Mol. Ecol. Resour.* 17, 1136–1147
- O'Connell, J., Schulz-Trieglaff, O., Carlson, E., Hims, M.M., Gormley, N.A. & Cox, A.J. (2015). NxTrim: optimized trimming of Illumina mate pair reads. *Bioinformatics*, 31, 2035–2037.
- Orr, H.A. (2005). The genetic theory of adaptation: a brief history. *Nat. Rev. Genet.*, 6, 119–127.
- Pacifici, M., Foden, W.B., Visconti, P., Watson, J.E., Butchart, S.H., Kovacs, K.M. *et al.* (2015). Assessing species vulnerability to climate change. *Nat. Clim. Chang.*, 5, 215–224.
- Pardieck, K.L., Ziolkowski, D.J. Jr., Lutmerding, M., Campbell, K. & Hudson, M.-A.R. (2017). North American breeding bird survey. *J. Fish Wildlife Manag.* 8, 154–172
- Parmesan, C. & Yohe, G. (2003). A globally coherent fingerprint of climate change impacts across natural systems. *Nature*, 421, 37–42.
- Paxton, E.H. (2000). *Molecular Genetic Structuring and Demographic History of the Willow Flycatcher (Empidonax traillii)*. Northern Arizona University Flagstaff, AZ, USA.
- Rimoldi, S., Lasagna, E., Sarti, F.M., Marelli, S.P., Cozzi, M.C., Bernardini, G. *et al.* (2015). Expression profile of six stress-related genes and productive performances of fast and slow growing broiler strains reared under heat stress conditions. *Meta Gene*, 6, 17–25.
- Sauer, J.R., Pardieck, K.L., Ziolkowski, Jr D.J., Smith, A.C., Hudson, M.A., Rodriguez, V., Berlanga, H., Niven, D.K., Link, W.A. (2017). The first 50 years of the North American Breeding Bird Survey. *The Condor*, 119, 576–93.
- Savolainen, O., Lascoux, M. & Merilä, J. (2013). Ecological genomics of local adaptation. *Nat. Rev. Genet.*, 14, 807–820.
- Sedgwick, J.A. (2000). Willow flycatcher (*Empidonax traillii*). In: *The Birds of North America* (eds A.F. Poole and F.B. Gill). Cornell Lab of Ornithology Ithaca, NY, USA.
- Service, U.S.F.A.W. (2002). Final Recovery Plan Southwestern Willow Flycatcher (*Empidonax traillii* extimus). (ed. Service, USFaW) Albuquerque, New Mexico, p. 210.
- Sexton, J.O., Song, X.-P., Feng, M., Noojipady, P., Anand, A., Huang, C. *et al.* (2013). Global, 30-m resolution continuous fields of tree cover: Landsat-based rescaling of MODIS vegetation continuous fields with lidar-based estimates of error. *Inter. J. Digital Earth*, 6, 427–448.
- Sinervo, B., Mendez-De-La-Cruz, F., Miles, D.B., Heulin, B., Bastiaans, E., Villagrán-Santa Cruz, M. *et al.* (2010). Erosion of lizard diversity by climate change and altered thermal niches. *Science*, 328, 894–899.
- Sogge, M.K., Marshall, R.M., Sferra, S.J. & Tibbitts, T.J. (1997). A southwestern willow flycatcher natural history summary and survey protocol. Technical Report NPS/NAUCPRS/NRTR-97/12. USGS Colorado Plateau Research Station, Northern Arizona University, Flagstaff, AZ.
- Stein, L. (2001). Genome annotation: from sequence to biology. *Nat. Rev. Genet.*, 2, 493.
- Stephens, P.A., Mason, L.R., Green, R.E., Gregory, R.D., Sauer, J.R., Alison, J. *et al.* (2016). Consistent response of bird populations to climate change on two continents. *Science*, 352, 84–87.
- Tarailo-Graovac, M. & Chen, N. (2009). Using RepeatMasker to identify repetitive elements in genomic sequences. *Current protocols in bioinformatics*, 5, 4–10.
- Theimer, T.C., Smith, A.D., Mahoney, S.M. & Ironside, K.E. (2016). Available data support protection of the Southwestern Willow Flycatcher Under the Endangered Species Act. *Condor*, 118, 289–299.
- Unitt, P. (1987). *Empidonax traillii* extimus: an endangered subspecies. *West. Birds*, 18, 137–162.
- Urban, M.C. (2015). Accelerating extinction risk from climate change. *Science*, 348, 571–573.
- Warren, R., VanDerWal, J., Price, J., Welbergen, J., Atkinson, I., Ramirez-Villegas, J. *et al.* (2013). Quantifying the benefit of early climate change mitigation in avoiding biodiversity loss. *Nat. Clim. Chang.*, 3, 678–682.

- Wiens, J.J. (2016). Climate-related local extinctions are already widespread among plant and animal species. *PLoS Biol.*, 14, e2001104.
- Williams, S.E., Shoo, L.P., Isaac, J.L., Hoffmann, A.A. & Langham, G. (2008). Towards an integrated framework for assessing the vulnerability of species to climate change. *PLoS Biol.*, 6, 2621–2626.
- Zdobnov, E.M. & Apweiler, R. (2001). InterProScan—an integration platform for the signature-recognition methods in InterPro. *Bioinformatics*, 17, 847–848.
- Zhang, J., Schmidt, C.J. & Lamont, S.J. (2017). Transcriptome analysis reveals potential mechanisms underlying differential heart development in fast-and slow-growing broilers under heat stress. *BMC Genom.*, 18, 295.
- Zheng, X., Levine, D., Shen, J., Gogarten, S.M., Laurie, C. & Weir, B.S. (2012). A high-performance computing toolset for relatedness and principal component analysis of SNP data. *Bioinformatics*, 28, 3326–3328.
- Zink, R.M. (2015). Genetics, morphology, and ecological niche modeling do not support the subspecies status of the endangered Willow Flycatcher (*Empidonax traillii extimus*). *Condor*, 117, 78–86.

## SUPPORTING INFORMATION

Additional Supporting Information may be found online in the supporting information tab for this article.

Editor, Tim Coulson

Manuscript received 9 November 2017

First decision made 11 December 2017

Second decision made 22 February 2018

Manuscript accepted 15 March 2018